

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 33/53, 33/531, C07K 17/00, C07H 1/00	A1	(11) International Publication Number: WO 95/35503 (43) International Publication Date: 28 December 1995 (28.12.95)
(21) International Application Number: PCT/US95/08882 (22) International Filing Date: 21 June 1995 (21.06.95) (30) Priority Data: 08/263,804 22 June 1994 (22.06.94) US (60) Parent Application or Grant (63) Related by Continuation US 08/263,804 (CIP) Filed on 22 June 1994 (22.06.94) (71) Applicant (for all designated States except US): PHARMA- COPEIA, INC. [US/US]; 101 College Road East, Princeton, NJ 08540 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BALDWIN, John, J. [US/US]; 621 Gypsy Hill Circle, Gwynedd Valley, PA 19437 (US). HORLBECK, Eric, G. [US/US]; 24-10 Ravens Crest Drive, Plainsboro, NJ 08536 (US). (74) Agents: LOPEZ, Gabriel et al.; Pharmacoepia, Inc., 101 College Road East, Princeton, NJ 08540 (US).	(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>	
(54) Title: SYNTHESIS OF COMBINATORIAL LIBRARIES (57) Abstract Directly dividing the contents of each sub-pool into the sub-pools for the next step in the synthetic scheme for producing a combinatorial library reduces the standard deviation, σ , relative to the standard deviation of the split synthesis method for producing such libraries.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

WO 95/35503

PCT/US95/08882

TITLE OF THE INVENTION

SYNTHESIS OF COMBINATORIAL LIBRARIES

BACKGROUND OF THE INVENTION

There is interest in methods for the synthesis of large numbers of diverse compounds which can be screened for various possible physiological or other activities. Techniques have been developed in which one adds individual units sequentially as part of the chemical synthesis to produce all or a substantial number of the possible compounds which can result from all the different choices possible at each sequential stage of the synthesis. Numerous strategies have been devised for producing such combinatorial libraries (R. M. Baum, *Combinatorial Approaches Provide Fresh Leads for Medicinal Chemistry*, C&EN, 20-25, Feb. 7, 1994). Common to many of these strategies is a technique known as split synthesis (Furka *et al.*, "Cornucopia of Peptides by Synthesis", Abstr. 14th Int. Cong. of Biochem., Prague Czechoslovakia, Vol. 5, p 47 (1988) and "More Peptides by Less Labour", Abstr. 10th Int. Sym. on Med. Chem., Budapest, Hungary p 288 (1988)), which essentially comprises dividing a first pool of material into sub-pools, treating these sub-pools so as to effect a change in the material, mixing the sub-pools into a second pool, and then again dividing the changed material into a new set of sub-pools for further treatment. This process is iterated until the desired end products are produced. Ideally, every theoretical member of the library is produced by this synthesis on solid supports, where each solid support holds an individual member, and these members are present in equal number, i.e., in the final pool there is an equal number of members for every possible combination of reactions. However, a perfectly even splitting is not achieved and this can lead to the non-production of some members and an over or under production of others. Over or under production of members can lead to other errors. For example, in deconvolution analysis of combinatorial libraries over or under representation of library members can be misinterpreted as indicating strong or weak compounds, e.g., in a binding assay, and result in the further elaboration of uninteresting compounds. One way

WO 95/35503

PCT/US95/08882

-2-

to reduce the first of these possibilities (non-production) is to design for an excess amount of each member in these syntheses, typically 10-1000 fold redundancy. However, this approach does not affect the over or under production of certain members. Even with redundancy there is a calculable error (σ) from the ideal. Reduction of this error is desirable.

SUMMARY OF THE INVENTION

It has now been found that directly dividing the contents of each sub-pool into the sub-pools for the next step in the synthetic scheme reduces σ relative to the σ of split synthesis.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a method for producing combinatorial libraries of compounds. These libraries of compounds can include 1) biopolymers such as peptides, oligonucleotides, and oligo-saccharides, 2) combinations of peptides, oligonucleotides, and oligo-saccharides, 3) non-oligomers such as heterocyclic, aromatics, alicyclics, aliphatics, and combinations thereof, and 4) any other compounds amenable to combinatorial synthesis. Non-oligomers includes benzodiazepines, prostaglandins, sulfonamides, pyrrolidines, etc. Combinatorial libraries of compounds are produced by partitioning a set of particles, i.e., the solid supports, into a first group of reaction vessels wherein they uniquely treated, re-partitioning the treated particles via a common pool into a second group of reaction vessels wherein they are again uniquely treated, and repeating this process until the desired combinatorial library is produced. The unique treatment in each vessel is typically the addition, as by chemical means, of some moiety, each such treatment being selected so that at the end of the process all the members of the combinatorial library will have been produced. The method of this invention, which is referred to herein as direct dividing, comprises directly dividing among the reaction vessels to be used for the next step, in substantially equal amounts, the particles in each reaction vessel at the end of the previous step. Unlike split synthesis, at no time during the synthesis are the treated particles combined into a common pool. In this way, each reaction vessel in a

WO 95/35503

PCT/US95/08882

-3-

next step will contain substantially the same number of identical particles from each reaction vessel in the previous step. The procedure may be repeated, typically 1-5 times, until the synthesis is complete. Preferably, at the end of the synthesis every member of the library is represented in equimolar amounts.

Directly dividing the reaction supports offers a significant advantage over split synthesis, i.e., pooling and dividing, since it distributes more evenly the different compounds created at each step. This means that the actual number of solid supports holding non-identical library members will be closer to the theoretical minimum number of supports. As a result there will be fewer cases of compounds not represented, over-represented, or under-represented in the final pool. Consequently, this reduces the redundancy necessary to assure a minimum expected representation of each possible compound in the library. For example, 10,000 beads are equally divided into 100 reaction vessels. A different bifunctional synthon is allowed to react with the solid supports in each reaction vessel. Under split synthesis, the solid supports are combined and then equally divided into another 100 reaction vessels where they are allowed to react with a second synthon. A combinatorial analysis of this approach shows that 36.4% of the possible products are not produced at all and 26.4% of the possible products are over-represented. If the number of beads is increased to 30,000 (i.e., 3-fold redundancy), the number of products not represented is reduced to 4.8% but there is still be a significant number of products over-represented (35.3%) and under-represented (42.1%). In contrast, using the direct division method of this invention, every member of the combinatorial library is produced in equal proportion using either 10,000 or 30,000 beads.

The structure of each compound on a given solid support in a combinatorial library is determined by its reaction history, i.e., by the identity of the reaction vessels for each of its starting materials and intermediates. Ideally, each possible compound would be represented in the final mixture in a pre-determined amount. This will deviate based on a statistical variation inherent to the splitting paradigm.. In order for the theoretical outcome to occur, at each reaction step, each reaction

WO 95/35503

PCT/US95/08882

-4-

vessel must contain a pre-determined amount of every compound produced up to that point. However, when a mixture of intermediates obtained by pooling each subdivision from the previous step is divided into the reaction vessels for the next step in the synthesis, there is a statistical variation in the intermediates which go into each vessel. Such variation is inherent for both split synthesis and direct division after the second synthetic step. However, the variation (σ) will be significantly less for direct dividing than for split synthesis.

Each time that splitting occurs, the probability that x members of a given compound will be drawn into a given vessel is described by the hypergeometric distribution,

$$P(x) = \frac{\binom{s}{x} \binom{n-s}{d-x}}{\binom{n}{d}},$$

wherein

- s is the number of members of the given compound in the original vessel,
 n is the total number of components (i.e., (compounds)•(redundancy)) in the original vessel, and
 d is the number of components drawn from the original vessel into the given vessel.

The expected number of members of a given compound drawn is $\bar{x} = \frac{ds}{n}$ (where \bar{x} is the mean of x). A measure of how much x varies about the mean is given by the variance,

$$\sigma^2 = \frac{d(n-d)s(n-s)}{n^2(n-1)}.$$

- The following analysis applies under these pre-conditions:
- 1) there are more than one reaction vessel at each of step i and $i + 1$;
 - 2) there are at least as many intermediates as vessels at step i ; and
 - 3) there are more than one intermediate that have been through the path of

WO 95/35503

PCT/US95/08882

-5-

interest up to step i . When a split occurs, if s_i is the number of intermediate compounds that have been through a particular synthetic path up and including step i , then $P(s_{i+1})$ is the probability distribution of those intermediates that continue on a particular path up to step $i+1$.

- 5 Both split synthesis and direct dividing will have such a distribution for apportioning the intermediates from step i to step $i+1$ given s_i intermediates of interest in step i . The difference will be the appropriate values of n and d .

Now let b_i = number of vessels in step i .

- 10 For split synthesis:

n = total number of beads in the mixing container = B , and

d = number of beads drawn into each container on level $i+1$ = $B/b_i + 1$.

This leads to a variance of

$$\sigma_{split}^2 = \frac{\frac{B}{b_{i+1}} \left(B - \frac{B}{b_{i+1}} \right) s(B-s)}{B^2 (B-1)} = \frac{\left(1 - \frac{1}{b_{i+1}} \right) s(B-s)}{b_{i+1} (B-1)}$$

- 15 For direct divide synthesis:

n = total number of beads in level i container = B/b_i , and

d = number of beads drawn from level $B/(b_i b_i + 1)$.

This leads to a variance of

$$\sigma_{direct}^2 = \frac{\frac{B}{b_i b_{i+1}} \left(\frac{B}{b_i} - \frac{B}{b_i b_{i+1}} \right) s \left(\frac{B}{b_i} - s \right)}{\left(\frac{B}{b_i} \right)^2 \left(\frac{B}{b_i} - 1 \right)} = \frac{\left(1 - \frac{1}{b_{i+1}} \right) s(B - b_i s)}{b_{i+1} (B - b_i)}$$

- 20 The smaller the variance, the more evenly distributed the different compounds will be on step $i+1$. To show which variance is smaller we take the difference,

WO 95/35503

PCT/US95/08882

-6-

$$\begin{aligned}\sigma_{split}^2 - \sigma_{direct}^2 &= \frac{\left(1 - \frac{1}{b_{i+1}}\right) s_i}{b_{i+1}} \left(\frac{B - s_i}{B - 1} - \frac{B - s_i b_i}{B - b_i} \right) \\ &= \frac{\left(1 - \frac{1}{b_{i+1}}\right) s_i}{b_{i+1}} \left(\frac{B(s_i - 1)(b_i - 1)}{(B - 1)(B - b_i)} \right)\end{aligned}$$

which is greater than zero if $b_i > 1, b_{i+1} > 1, s_i > 1, B > b_i$; i.e., the variance of direct dividing will always be smaller than the variance of split synthesis.

The present invention also relates to combinatorial chemical libraries of compounds, optionally encoded with tags, and to the use of these libraries in assays to discover biologically active compounds. The invention also relates to methods for their preparation, intermediates, and to methods and pharmaceutical formulations for using these compounds in the treatment of mammals, especially humans.

One embodiment of the invention is the use of a combinatorial library of compounds generated by the direct divide method in assays to discover biologically active compounds (ligands). Thus, an aspect of the invention is a method of identifying a compound having a desired characteristic which comprises synthesizing a combinatorial library and testing the compounds of this library and the ligands, either attached to the solid support or detached therefrom, in an assay which identifies compounds having the desired characteristic. A further embodiment of the invention is determining the structure of any compound so identified.

Definitions

The following abbreviations have the indicated meaning:

DEAD = diethylazodicarboxylate
DCM = dichloromethane = methylene chloride

WO 95/35503

PCT/US95/08882

-7-

	DIC	=	diisopropylcarbodiimide
	DMAP	=	4-N,N-dimethylaminopyridine
	DMF	=	N,N-dimethylformamide
	EDT	=	ethane dithiol
5	FACS	=	fluorescence activated cell sorting
	Fmoc	=	9-fluorenylmethoxycarbonyl
	HOBt	=	N-hydroxybenzotriazole
	Me	=	methyl
	PEG	=	polyethylene glycol
10	Ph	=	phenyl
	r.t.	=	room temperature
	sat'd	=	saturated
	t-	=	tertiary
	TFA	=	trifluoroacetic acid
15	THF	=	tetrahydrofuran

Any identification method may be used with this invention. Thus, a strategy using chemical entities, i.e., tags, or a deconvolution, or "mimotope strategy", method such as taught by Geysen *et al.*, J. Immunol Methods, **102**, 259-274 (1987) and Geysen *et al.*, Mol. Immunol. **23**, 709-715 (1986) may be used. The tags may be peptides (e.g., Kerr *et al.*, J. Amer. Chem. Soc., **115**, 2529-2531 (1993) and Nikolaiev *et al.*, Pept. Res., **6**, 161-170 (1993)), oligonucleotides (e.g., Needles *et al.*, *Proc. Natl. Acad. Sci. USA*, **90**, 10700-10704 (1993), Lerner *et al.*, PCT Appli. WO 93/20242, and Brenner and Lerner *Proc. Natl. Acad. Sci. USA*, **89**, 5181-5183 (1992)), or electrophoric tags (e.g., Ohlmeyer *et al.*, *Proc. Natl. Acad. Sci. USA*, **90**, 10922-10926 (1993) and Still *et al.*, PCT Appli. WO 94/08051). Preferably, the electrophoric tags are chemical entities which possess several properties: they must be detachable from the solid supports, preferably by photolysis or oxidation; they must be individually differentiable, and preferably separable; they must be stable under the synthetic conditions; they must be capable of being detected at very low concentrations, e.g., 10^{-18} to 10^{-9} mole; they should be identifiable with readily-available equipment which does not require sophisticated technical capabilities to operate; and they should be relatively

WO 95/35503

PCT/US95/08882

-8-

economical. The tags may be structurally related or unrelated, e.g., a homologous series, repetitive functional groups, related members of the Periodic Chart, different isotopes, combinations thereof, and the like. At the end of the combinatorial synthesis, to each solid support, there
5 will usually be attached at least 0.01 femtomol, usually 0.001-50 pmol, of each tag. The tags may be aliphatic, alicyclic, aromatic, heterocyclic, or combinations thereof. Distinguishing features may be the number of repetitive units, such as methylene groups in an alkyl moiety; alkyleneoxy groups in a polyalkyleneoxy moiety; halo groups in a
10 polyhalo compound; α - and/or β -substituted ethylene groups where the substituents may be alkyl groups, oxy, carboxy, amino, halo, or the like; isotopes; etc.

The materials upon which the combinatorial syntheses of the invention are performed are referred to as solid supports, beads,
15 and resins. These terms are intended to include:

beads, pellets, disks, fibers, gels, or particles such as cellulose beads, controlled pore-glass beads, silica gels, polystyrene beads optionally cross-linked with
divinylbenzene and optionally grafted with polyethylene
20 glycol and optionally functionalized with amino, hydroxy, carboxy, or halo groups, grafted co-poly beads, polyacrylamide beads, latex beads, dimethylacrylamide beads optionally cross-linked with N,N'-bis-acryloyl ethylene diamine, and glass particles coated with hydrophobic
25 polymer, etc., i.e., material having a rigid or semi-rigid surface.

Optical Isomers - Diastereomers - Geometric Isomers

Some of the compounds described herein contain one or more asymmetric centers and may thus give rise to enantiomers,
30 diastereomers, and other stereoisomeric forms which may be defined in terms of absolute stereochemistry as (R) or (S), or as (D) or (L) for amino acids. The present invention is meant to comprehend all such possible diastereomers as well as their racemic and optically pure forms. Optically active (R) and (S), or (D and L), isomers may be
35 prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. When the compounds described herein contain

WO 95/35503

PCT/US95/08882

-9-

olefinic double bonds, and unless specified otherwise, it is intended to include both E and Z geometric isomers.

Utilities

These compounds may also be used as libraries for


5 discovering new lead structures by evaluation across an array of biological assays, including the discovery of selective inhibition patterns across isozymes. These libraries are thus tools for drug discovery; i.e., as a means to discover novel lead compounds by screening the libraries against a variety of biological targets and to develop structure-activity

10 relationships in large families of related compounds. The libraries may be tested with the ligands attached to the solid supports or the individual compounds may be detached prior to evaluation. With the compounds of the combinatorial library, screening assays such as FACS sorting and cell lawn assays may be used. When a ligand is detached prior to

15 evaluation, its relationship to its solid support is maintained, for example, by location within the grid of a standard 96-well plate or by location of activity on a lawn of cells. The solid support associated with bioactivity or the solid support related to the detached ligand may then be decoded to reveal the structural or synthetic history of the active

20 compound (Ohlmeyer *et al.*, *Proc. Natl. Acad. Sci. USA*, 90, 10922-10926, Dec. 1993).

The following schemes demonstrate application of the method of the invention to the preparation of a tagged combinatorial library of sulfonamides represented by Formula I:



wherein:


\textcircled{S} is a solid support;

T'-L- is of the Formula:

WO 95/35503

PCT/US95/08882

-10-




wherein $n = 3-12$ when Ar is pentachlorophenyl and
 $n = 4-6$ when Ar is 2,4,6-trichlorophenyl;

5

q is 3-13;
 -L'-II' is a ligand/linker residue;
 -L'- is

10



15

wherein the left-hand bond as shown is the point of attachment to the solid support and the right hand bond is the point of attachment to the ligand, and B is O or NH, with the proviso that in (b) B is NH when attached to a carbonyl group in the ligand; and

20

II is represented by the formula



wherein R^1 , A, and Y are as defined in Tables 1-1, 1-2, and 1-3, respectively.

25

30

At each step in the synthesis each solid support upon which a compound is being synthesized is uniquely tagged to define the particular chemical event(s) occurring during that step. The tagging is accomplished using identifiers such as those of Formula IV, which record the sequential events to which the support is exposed during the synthesis, thus providing a reaction history for the compound produced on each support. The identifiers are used in combination with one another to form a binary or higher order encoding scheme permitting a relatively small number of identifiers to encode a relatively large number of reaction products.

WO 95/35503

PCT/US95/08882

-11-

In carrying out the synthesis, one begins with at least 4×10^4 , and generally not exceeding 10^{15} solid supports. Depending on the pre-determined number of R^1 choices for the first step, one divides the supports accordingly into as many containers. The appropriate reagents and reaction conditions are applied to each container and the combination of identifiers which encode for each R^1 choice is added and attached. Depending on the chemistries involved, the tagging may be done prior to, concomitantly with, or after the reactions which comprise each choice. As a control, sample supports may be picked at any stage and a portion of their tags detached and decoded to verify that the correct tags are bound to the sample supports. As needed, one may wash the beads free of any excess reagents or by-products before proceeding. At the end of each step, the supports in each container are divided into as many containers as pre-determined for the number of A choices for the second step in the synthesis. Direct dividing is accomplished by, e.g., weighing or by the isopycnic method of pipetting aliquots from an isobuoyant suspension of supports. This procedure of dividing, reacting, tagging, and direct dividing is repeated until the combinatorial synthesis is completed.

Scheme 1

Functionalized supports such as amino-functionalized or hydroxy-terminating PEG grafted polystyrene beads are equally divided into a pre-determined number of reaction vessels and are reacted with either a) a cleavable linker/ligand element which has been pre-formed to generate 2 when the detached ligand element terminates in OH or b) a cleavable linker, followed by reaction with a ligand element to generate 3 in the case when the detached ligand element terminates in COOH. Compounds 2 and 3 are then treated with piperidine/DMF to de-protect the amino group of the ligand element R^1 to yield 4 (Scheme 1, step (c)). Unique tagging of the supports in each reaction vessel is achieved with combinations of identifiers encoded in a binary scheme, e.g., as depicted in Table 1-1 for seven choices of R^1 . The identifiers are attached by adding a solution of the identifiers (in a 5% wt./wt. identifier:solid support ratio) to a batch of supports suspended in CH_2Cl_2 and shaking the mixture for 30 min. A dilute solution of

WO 95/35503

PCT/US95/08882

-12-

rhodium trifluoroacetate dimer is added and the mixture is immediately shaken. The mixture is continued to be shaken overnight and then washed in CH₂Cl₂.

Scheme 2

- 5 The compounds 4 are directly divided into a pre-determined number of reaction vessels, each of which is treated with one reagent corresponding to ligand element A, in the presence of HOBt/DMF to produce 5. The reaction vessels are then treated with piperidine/DMF to de-protect the terminal amino group, yielding 6.
- 10 Unique tagging of the supports in each reaction vessel is achieved with combinations of additional identifiers encoded in a binary scheme, e.g., as depicted in Table 1-2 for 31 choices of A.

Scheme 3

- 15 The compounds 6 are then directly divided into a pre-determined number of reaction vessels each of which is treated with one reagent corresponding to ligand element Y in the presence of solvents such as CH₂Cl₂ and DMF, and, when required, condensation reagents such as DIC, acylation catalysts such as DMAP, and bases such as triethylamine to produce 7. Unique tagging of the supports in each
- 20 reaction vessel is achieved with combinations of additional identifiers encoded in a binary scheme, e.g., as in Table 1-3 for 31 choices of Y. Compound 7 is then either exposed to UV light (~360 nm) in a lower alkanol such as MeOH to cleave the protected form of the compounds of
- 25 Formula II from the support/linker complex or first treated with TFA/thioanisole/EDT to remove the protecting groups on the R² sidechains and then exposed to UV light in a lower alkanol such as MeOH to cleave compound II.



WO 95/35503

PCT/US95/08882





-13-

SCHEME 1 ADDITION OF R¹

a. Carbonate-linked Moieties



Reagents 1-4




WO 95/35503

PCT/US95/08882


-14-

SCHEME 1 (Cont.)

b. Ester-linked Moieties



c. Fmoc Deprotection



* where R¹ includes the
carbonate linkage in 2 and
the ester linkage in 3


30

WO 95/35503

PCT/US95/08882

-15-

SCHEME 2
ADDITION OF A




(Prot) = optional, non-Fmoc, base stable sidechain protecting group

WO 95/35503

PCT/US95/08882

-16-

SCHEME 3
ADDITION OF Y



WO 95/35503

PCT/US95/08882


-17-

The invention is further defined by reference to the following examples, which are intended to be illustrative and not limiting.

PREPARATION 1

IDENTIFIERS

Thirteen compounds of the general formula:



wherein:

- 10 $n = 3-12$ and Ar is pentachlorophenyl or
 $n = 4-6$ and Ar is 2,4,6-trichlorophenyl
 were prepared according to Scheme 4 and the following illustrative
 example.

- 15 a) Methyl vanillate (0.729 g, 4.0 mmole), 1-hydroxy-9-(2,3,4,5,6-pentachlorophenoxy)nonane (1.634 g, 4.0 mmole) and triphenylphosphine (1.258 g, 4.8 mmole) were dissolved in 20 mL dry toluene under argon. DEAD (0.76 mL, 0.836 g, 4.8 mmole) was added dropwise and the mixture was stirred at 25°C for one hr. The solution was concentrated to half volume and purified by flash chromatography
 20 eluting with DMC to give 1.0 g (1.7 mmole, 43%) of the product as a white crystalline solid.

- 25 b) The methyl ester from Step (a) (1.0 g, 1.7 mmole) was dissolved in 50 mL THF, 2 mL water was added, followed by LiOH (1.2 g, 50 mmole). The mixture was stirred at 25°C for one hr. then refluxed for 5 hr. After cooling to 25°C, the mixture was poured onto ethyl acetate (200 mL) and the solution was washed with 1 M HCl (3x 50 mL) then sat'd aq. NaCl (1x 50 mL) and dried over sodium sulfate. The solvent was removed and the crude acid azeotroped once with toluene.

- 30 c) The crude material from Step (b) was dissolved in 100 mL toluene, 10 mL (1.63 g, 14 mmole) thionyl chloride was added,

WO 95/35503

PCT/US95/08882

-18-

- and the mixture was refluxed for 90 min. The volume of the solution was reduced to approx. 30 mL by distillation, then the remaining toluene was removed by evaporation. The crude acid chloride was dissolved in 20 mL dry DCM and cooled to -70°C under argon and a
- 5 solution of approx. 10 mmole diazomethane in 50 mL anhydrous ether was added. The mixture was warmed to r.t. and stirred for 90 min. Argon was bubbled through the solution for 10 min., then the solvents were removed by evaporation and the crude material was purified by flash chromatography, eluting with 10-20% ethyl acetate in hexane.
- 10 The diazoketone (0.85 g, 1.4 mmole, 82% yield over three steps) was obtained as a pale yellow solid.

The other 12 identifiers of Formula IV were prepared by analogous synthetic routes, steps (a), (b), and (c).

- In the synthesis of Example 1, the 13 identifiers were used
- 15 to encode the combinatorial library. In Step 1, pentachlorophenyl identifiers where $n = 10-12$ (abbreviated $C_{10}Cl_5$, $C_{11}Cl_5$, and $C_{12}Cl_5$) were used in the following binary encoding scheme: 001 = ($n = 12$), 010 = ($n = 11$) and 100 = ($n = 10$). In Step 2, pentachlorophenyl identifiers where $n = 5-9$ (abbreviated C_5Cl_5 , C_6Cl_5 , C_7Cl_5 , C_8Cl_5 , and
- 20 C_9Cl_5) were used and encoded as follows: 00001 = ($n = 9$), 00010 = ($n = 8$), 00100 = ($n = 7$), 01000 = ($n = 6$) and 10000 = ($n = 5$). In Step 3, pentachlorophenyl identifiers where $n = 3-4$ (abbreviated C_3Cl_5 and C_4Cl_5) were used and encoded as follows: 00001 = ($n = 4$), 00010 = ($n = 3$). Also in Step 3, trichlorophenyl identifiers where $n = 4-6$
- 25 (abbreviated C_4Cl_3 , C_5Cl_3 , and C_6Cl_3) were used and encoded as follows: 00100 = ($n = 6$), 01000 = ($n = 5$), and 10000 = ($n = 4$).


- Thus, in Step 1 reagent 3 (Table 1-1) is encoded "011" which represents tagging this choice in the synthesis with the two pentachloro-phenyl identifiers where $n = 11$ and 12. Likewise, in Step
- 30 3 reagent 52 (Table 1-3) is encoded "01110" which represents tagging this choice in the synthesis with the pentachlorophenyl identifier where $n = 3$ and the two trichlorophenyl identifiers where $n = 5$ and 6.

WO 95/35503

PCT/US95/08882

-19-

SCHEME 4
IDENTIFIERS



WO 95/35503

PCT/US95/08882

-20-

PREPARATION 2
t-BUTYL 4-(HYDROXYMETHYL)-3-NITROBENZOATE

t-Butyl 4-(acetoxymethyl)-3-nitrobenzoate was prepared as described by Barany and Albericio, *J. Am. Chem. Soc.* **1985**, *107*, 4936-4942. However, the reference's final procedure for hydrazinolysis of the acetate using hydrazine hydrate in CHCl_3 at 25°C to produce t-butyl 4-(hydroxymethyl)-3-nitrobenzoate, which is the t-butyl ester pre-cursor of the photocleavable linker used herein, was changed to hydrazinolysis using hydrazine hydrate in MeOH at 25°C to produce the desired hydroxymethyl final product in near quantitative yield.

This new hydrazinolysis procedure was also used to produce the photocleavable linker when attached to TentaGel S NH_2 resin.

t-Butyl 4-(hydroxymethyl)-3-nitrobenzoate: To a solution of 14.1 g (47.7 mmol, 1.00 eq.) of t-butyl 4-(acetoxymethyl)-3-nitrobenzoate in MeOH (200 mL) was added 27.0 mL (477 mmol, 10.0 eq.) of hydrazine hydrate (55% hydrazine). The resulting yellow solution was stirred at 25°C for 4 hr. EtOAc (250 mL) and saturated aq. NaCl (85 mL) were added, and the organic layer collected after shaking. The organic layer was washed further with saturated aq. NaCl (2 x 85 mL), and then dried (MgSO_4). Removal of volatiles *in vacuo* gave the product in 93% yield as yellow crystals.

TentaGel S NH_2 - attached 4-(hydroxymethyl)-3-nitrobenzamide: See Example 1, Step 1a.


The reaction is represented in Scheme 5.

WO 95/35503

PCT/US95/08882

-21-

SCHEME 5 HYDRAZINOLYSIS



EXAMPLE 1

6727 COMPOUND LIBRARY

Step 1

a) Preparation of ester-linked resin batches

20 TentaGel S NH₂, (10 g, 2.9 mmole, 0.29 mmole/g) was suspended in 60 mL methylenechloride. 4-acetoxymethyl-3-nitrobenzoic acid (2.77 g, 11.6 mmole, 4 eq), DMAP (1.42 g, 11.6 mmole, 4 eq) and DIC (1.81 mL, 1.46 g, 11.6 mmole, 4 eq) was added in that order. The mixture was agitated at 25 °C with a wrist action shaker for 5 hours, at which time the resin gave a negative Kaiser test. The derivatized resin was washed (methylenechloride 5 x 50 mL, isopropanol 5 x 50 mL, then methylenechloride 5 x 50 mL), dried *in vacuo* and stored in the dark.

25 The acetoxymethyl resin above was washed with methanol (1 x 150 mL), then suspended in 150 mL 10% hydrazine/methanol. The resin was agitated at 25 °C for 44 hours, filtered, then washed (methanol, 5 x 100 mL, then methylene chloride, 5 x 100 mL) and dried *in vacuo*.

30 Three batches of the above hydroxymethyl resin (3 g each, 0.84 mmole, 0.28 mmole/g) were placed in 100 mL synthesis vessels and each was suspended in 60 mL dichloromethane. One of N-Fmoc isonipecotic acid (0.885 g, 2.52 mmole, 3 eq), N-Fmoc nipecotic acid

WO 95/35503

PCT/US95/08882

-22-

(0.885 g, 2.52 mmole, 3 eq), and N-Fmoc- ϵ -amino caproic acid (0.890 g, 2.52 mmole, 3 eq) was added to each of the three batches. DMAP (0.031 g, 0.252 mmole, 0.3 eq) and then DIC (0.4 mL, 0.318 g, 2.52 mmole, 3 eq) was added to each and the resin batches were agitated at 25 °C for 22 hours. The resin batches were filtered, washed (methylenechloride (5 x 50 mL) and isopropanol (2 x 50 mL), then methylenechloride (5 x 50 mL)), and dried *in vacuo*.

b) Preparation of carbonate-linked resin batches

N-Fmoc-2-hydroxymethylpiperidine (1.35 g, 4 mmole, 1 eq) was dissolved in 20 mL methylenechloride and cooled to 0 °C. Phosgene (8 mL, 2 M solution in toluene, 16 mmole, 4 eq) was added followed by 2,6-lutidine (1.86 mL, 1.71 g, 4 eq), then the mixture was stirred for 30 min. The mixture was concentrated on a rotary evaporator to give a viscous slurry and this residue was redissolved in 30 mL methylenechloride (the solution contained some undissolved solid). ^tButyl 4-hydroxymethyl-3-nitrobenzoate (0.51 g, 2 mmole, 0.5 eq) was added and the mixture was stirred at 25 °C for 2 hours. The crude reaction mixture was poured onto 200 mL ethyl acetate and this solution was washed with 1 M HCl (2 x 100 mL), sat'd aq. NaHCO₃ (2 x 100 mL) and sat'd aq. NaCl (1 x 100 mL). The solution was dried over magnesium sulfate, filtered, and evaporated to give the crude product. This was purified by flash chromatography eluting with 10-30% ethylacetate/hexane to give the product (1.22 g, 1.97 mmole, 99%) as a pale yellow solid.

The carbonate above was dissolved in 20 mL methylene chloride, 10 mL TFA was added and the mixture was stirred at 25 °C for 16 hours. The solution was diluted with 50 mL toluene and evaporated to give the carboxylic acid as a viscous yellow oil which was azeotroped with toluene once, then dried *in vacuo*.

The acid prepared above (~2 mmole, 2 eq) was dissolved in 30 mL methylenechloride and this solution was added to TentaGel S NH₂ (3 g, 0.32 mmole/g, ~1 mmole, 1 eq). The resin was suspended by agitation then DMAP (40 mg, 0.3 mmole, 0.3 eq) and DIC (0.47 mL, 0.4 g, 3 eq) were added in that order. The resin was agitated at 25 °C for 19 hours, at which time it gave a negative Kaiser test. The resin

WO 95/35503

PCT/US95/08882

-23-

was filtered and washed (methylenechloride 10 x 50 mL), then dried *in vacuo*.

The three other carbonate linked resin batches were prepared in an analogous manner using the reagents of Table 1-1.

5 c) Encoding of Step 1

One gram batches of the seven resin batches from Steps 1(a) and 1(b) with the appropriately linked N-Fmoc protected amino acid or amino alcohol were placed in seven separate synthesis vessels and each was suspended in 20 mL methylene chloride.

- 10 Stock solutions of 200 mg of C₁₂Cl₅, C₁₁Cl₅, and C₁₀Cl₅-linker-diazoketone (Preparation 1) were prepared in 4 mL methylenechloride. Aliquots (1 mL) of the stock solutions were used to prepare the appropriate binary coding mixtures for each of the seven resin batches. The appropriate coding mixture was added to each resin
15 batch and the resin was agitated for 1 hour. Rhodium trifluoroacetate dimer (1 mL of a 1 mg/mL solution in methylenechloride) was added to each of the vessels and the resin was agitated at 25 °C overnight. Each resin batch was washed with methylenechloride (1 x 50 mL), then the batches were combined and the entire library (seven compounds) was
20 washed with methylenechloride (10 x 50 mL).

d) Deprotection

- N-Fmoc protecting groups were removed by washing the resin once with DMF, filtering, then suspending the resin in 60 mL 50% piperidine/DMF and agitating at room temperature for 30 min. The
25 resin was filtered, washed (DMF (5 x 50 mL) and methylenechloride (10 x 50 mL)) and dried *in vacuo*.

Step 2

a) Addition of A

- 30 The dried resin from Step 1(d) was divided into 31 batches of 210 mg (~0.07 mmole), each of which was placed in a 20 mL synthesis vessel. Each of the reagents (0.25 g, > 0.4 mmole, > 6 eq) (Table 1-2) used in the second step of the synthesis (N-Fmoc amino acid with acid labile side chain protection where appropriate) was dissolved in 10 mL
35 methylenechloride. HOBt (1 mL of 1 mg/mL in DMF) was added and

WO 95/35503

PCT/US95/08882

-24-

the solutions were shaken briefly. Further DMF was added to those amino acids that had not completely dissolved. Each reagent solution was added to one of the 31 resin batches. DIC (0.2 mL, ~1 mmole) was added to each vessel and the resin was agitated at 25 °C overnight.

- 5 Each of the resin batches was filtered and washed separately (DMF (1 x 15 mL) and methylenechloride (10 x 15 mL)). The resin was suspended in 10 mL methylenechloride.

b) Encoding of Step 2

- 10 Stock solutions of 160 mg of C₉Cl₅, C₈Cl₅, C₇Cl₅, C₆Cl₅, and C₅Cl₅ -linker-diazoketone (Preparation 1) were prepared in 16 mL methylenechloride. Aliquots (1 mL) of the stock solutions were used to prepare the appropriate binary coding mixtures for each of the 31 resin batches (Table 1-2). The appropriate coding mixture was added to each of the resin batches and the resin was agitated for 30 min. Rhodium
15 trifluoroacetate dimer (1 mL of a 1 mg/mL solution in methylenechloride) was added to each of the vessels and the resin was agitated at 25 °C overnight. Each resin batch was filtered then washed separately with methylenechloride (1 x 15 mL). The batches were combined and the entire library (217 compounds) was washed with
20 methylenechloride (5 x 50 mL).

c) Deprotection

- N-Fmoc protecting groups were removed by washing the resin once with DMF, filtering, then suspending the resin in 60 mL 50% piperidine/DMF and agitating at room temperature for 30 min. The
25 resin was filtered, washed (DMF (5 x 50 mL) and methylenechloride (10 x 50 mL)) and dried *in vacuo*.

Step 3

a) Addition of Y

- 30 The dried resin from Step 2 (c) was divided into 31 batches of 150 mg (~0.05 mmole), each of which was placed in a 20 mL synthesis vessel. Each of the reagents (0.25 g, > 0.4 mmole, > 8 eq) used in the third step of the synthesis was dissolved in methylenechloride or DMF or a mixture of the two as appropriate (Table 1-3). Each reagent
35 solution was added to one of the 31 resin batches, then any coreagents

WO 95/35503

PCT/US95/08882

-25-

were added as required (see table). The resin batches were agitated at 25 °C overnight, then each was washed (DMF (1 x 10 mL) and methylenechloride (10 x 10 mL)).

b) Encoding of Step 3

- 5 Stock solutions of 200 mg of C₄Cl₅ and C₃Cl₅ -linker-diazoketone (Preparation 1) were prepared in 16 mL methylenechloride. Stock solutions of 600 mg of C₆Cl₃, C₅Cl₃, and C₄Cl₃ -linker-diazoketone (Preparation 1) were prepared in 16 mL methylenechloride. Aliquots (1 mL) of the stock solutions were used to
- 10 prepare the appropriate binary coding mixtures for each of the 31 resin batches (Table 1-3). The appropriate coding mixture was added to each of the resin batches and the resin was agitated for 30 min. Rhodium trifluoroacetate dimer (1 mL of a 1 mg/mL solution in methylenechloride) was added to each of the vessels and the resin was
- 15 agitated at 25 °C overnight. Each resin batch was filtered then washed separately with methylenechloride (1 x 15 mL). The batches were combined and the entire library (6727 compounds) was washed with methylenechloride (5 x 50 mL), then dried *in vacuo*.

c) Deprotection








- 20 The combined resins from Step 3(b) (~ 2 g) were suspended in 60 mL TFA/thioanisole/EDT (50/50/5) and shaken at room temperature overnight. The resins were filtered, washed (methylenechloride 10 x 50 mL), and dried *in vacuo*.

WO 95/35503

PCT/US95/08882

-26-

Table 1-1
R¹ Residues and Encoding Scheme

	R ¹ Residue	Binary Code
5	1. 	001
10	2. 	010
15	3. 	011
20	4. 	100
25	5. 	101
30	6. 	110
	7. 	111








WO 95/35503

PCT/US95/08882

-27-

Table 1-2

A Reagents and Encoding Scheme









	Reagent	Binary Code
5	8.  (L)	00001
10	9.  (D)	00010
15	10.  (D)	00011
20	11.  (L)	00100
	12.  (L)	00101
25	13.  (L)	00110
30	14.  (D)	00111

WO 95/35503

PCT/US95/08882

-28-

Table 1-2 (Cont.)









5	15.		01000
	16.		01001
10	17.		01010
15	18.		01011
20	19.		01100
25	20.		01101
	21.		01110
30	22.		01111

WO 95/35503

PCT/US95/08882

-29-

Table 1-2 (Cont.)









5	23.	 (D)	10000
10	24.	 (D)	10001
15	25.	 (L)	10010
20	26.	 (L)	10011
25	27.	 (D)	10100
30	28.	 (L)	10101
	29.	 (L)	10110
	30.	 (L)	10111

WO 95/35503

PCT/US95/08882

-30-

Table 1-2 (Cont.)







5	31.		11000
	32.		11001
10	33.		11010
15	34.		11011
20	35.		11100
	36.		11101
25	37.		11110
30	38.		11111

WO 95/35503

PCT/US95/08882

-31-

Table 1-3
Y Reagents and Encoding Scheme








	Y Reagent	Solvent	Co-reagent(s)	Binary Code
5				
10	39. 	DMF	0.3 mL DIC, 100 mg DMAP	00001
15	40. 	DMF	0.3 mL DIC, 100 mg DMAP	00010
20	41. 	DMF	0.3 mL DIC, 100 mg DMAP	00011
25	42. 	CH ₂ Cl ₂	0.5 mL NEt ₃ , 100 mg DMAP	00100
30	43. 	CH ₂ Cl ₂	0.5 mL NEt ₃ , 100 mg DMAP	00101
	44. 	CH ₂ Cl ₂	0.5 mL NEt ₃ , 100 mg DMAP	00110

WO 95/35503

PCT/US95/08882

-32-

Table 1-3 (Cont.)








5	45.		CH ₂ Cl ₂	0.5 mL NEt ₃ , 100 mg DMAP	00111
10	46.		CH ₂ Cl ₂	0.5 mL NEt ₃ , 100 mg DMAP	01000
15	47.		CH ₂ Cl ₂ /DMF (1:1)	0.5 mL NEt ₃ , 100 mg DMAP	01001
20	48.	n-BuSO ₂ Cl	CH ₂ Cl ₂	0.5 mL NEt ₃ , 100 mg DMAP	01010
25	49.		CH ₂ Cl ₂	0.5 mL NEt ₃ , 100 mg DMAP	01011
30	50.		CH ₂ Cl ₂	0.5 mL NEt ₃ , 100 mg DMAP	01100
	51.		CH ₂ Cl ₂	0.5 mL NEt ₃ , 100 mg DMAP	01101
	52.		CH ₂ Cl ₂	0.3 mL DIC, 100 mg DMAP	01110

WO 95/35503

PCT/US95/06882

-33-

Table 1-3 (Cont.)

5	53.		CH ₂ Cl ₂ /DMF (1:1)	0.3 mL DIC, 100 mg DMAP	01111
	54.		CH ₂ Cl ₂ /DMF (1:1)	0.3 mL DIC, 100 mg DMAP	10000
10	55.		CH ₂ Cl ₂	100 mg DMAP	10001
15	56.		CH ₂ Cl ₂ /DMF (1:1)	0.3 mL DIC, 100 mg DMAP	10010
	57.	MeO(CH ₂) ₅ -CO ₂ H	CH ₂ Cl ₂	0.3 mL DIC, 100 mg DMAP	10011
20	58.		CH ₂ Cl ₂	0.3 mL DIC, 100 mg DMAP	10100
	59.	H ₃ C-CH ₂ -CO ₂ H	CH ₂ Cl ₂	0.3 mL DIC, 100 mg DMAP	10101
25	60.		CH ₂ Cl ₂ /DMF (1:1)	0.3 mL DIC, 100 mg DMAP	10110
30	61.		CH ₂ Cl ₂ /DMF (1:1)	0.3 mL DIC, 100 mg DMAP	10111

WO 95/35503

PCT/US95/08882

-34-

Table 1-3 (Cont.)

5	62.		CH ₂ Cl ₂ /DMF (1:1)	0.3 mL DIC, 100 mg DMAP	11000
	63.		CH ₂ Cl ₂		11001
10	64.		CH ₂ Cl ₂		11010
15	65.		CH ₂ Cl ₂	0.5 mL NEt ₃	11011
	66.		CH ₂ Cl ₂		11100
20	67.		Fmoc-G-OH, DIC, DMAP, DMF, then piperidine/ DMF	2-Napthalene- sulfonyl chloride, NEt ₃ , DMAP, CH ₂ Cl ₂	11101
25	68.		Fmoc-G-OH, DIC, DMAP, DMF, then piperidine/ DMF	4-Biphenyl- carboxylic acid, DIC, DMAP, CH ₂ Cl ₂ /DMF	11110
30	69.		Fmoc-G-OH, DIC, DMAP, DMF, then piperidine/ DMF	Imidazole- acetic acid, DIC, DMAP, CH ₂ Cl ₂ /DMF	11111

WO 95/35503

PCT/US95/08882

-35-

EXAMPLE 26727 COMPOUND LIBRARY

5 The procedure of Example 1 is followed except that in Steps 1(c) and 2(b) the batches are not combined. Instead, each batch is individually deprotected and then the batches are directly divided into the appropriate number of batches for the next step.

EXAMPLE 3EXAMPLE 1 IN 5-FOLD REDUNDANCY

10 The procedure of Example 1 is followed except that a total of 33,635 solid supports are used.

EXAMPLE 4EXAMPLE 2 IN 5-FOLD REDUNDANCY

The procedure of Example 2 is followed except that a total of 33,635 solid supports are used.

15

EXAMPLE 5COMPARISON OF EXAMPLES 3 AND 4

Combinatorial analyses of Examples 3 and 4 produce the following results:

% Not Represented		% Under-represented		% Over-represented	
Split Syn.	Direct Div.	Split Syn.	Direct Div.	Split Syn.	Direct Div.
0.657	0.413	44.0	42.4	38.4	38.3

20

All publications and patent applications cited herein are incorporated by reference.

WO 95/35503

PCT/US95/08882

-36-

WHAT IS CLAIMED IS:

1. In a process for producing a combinatorial library of compounds which comprises 1) partitioning a set of particles into a first group of reaction vessels wherein they are uniquely treated, 2) re-partitioning the treated particles via a common pool into a second group of reaction vessels wherein they are again uniquely treated, and 3) optionally, repeating step (2), the improvement which comprises re-partitioning the treated particles by directly dividing the contents of each reaction vessel into the reaction vessels for the next step in the synthesis.
2. A process of Claim 1 wherein the combinatorial library comprises peptides, oligonucleotides, or oligo-saccharides, or combinations thereof.
3. A process of Claim 1 wherein the combinatorial library comprises non-oligomers.
4. A process of Claim 3 wherein the non-oligomers are heterocyclic, aromatics, alicyclics, aliphatics, or combinations thereof.
5. A process of Claim 1 wherein the combinatorial library comprises compounds encoded with peptide or oligonucleotide tags.
6. A process of Claim 1 wherein the combinatorial library comprises compounds encoded with electrophoric tags.
7. A process of Claim 1 wherein each member is represented in equimolar amounts.
8. A process of Claim 1 wherein step (2) is repeated 1-5 times.
9. A pharmaceutical composition comprising a therapeutically effective amount of a compound produced by the process of Claim 1 and a pharmaceutically acceptable carrier.
10. A method of identifying a ligand having a desired characteristic which comprises synthesizing a combinatorial library by

WO 95/35503

PCT/US95/08882

-37-

the process of Claim 1 and testing the compounds in said library in an assay which identifies compounds having the desired characteristic.

11. A method of identifying a ligand having a desired characteristic which comprises synthesizing a combinatorial library by
5 the process of Claim 1, detaching the ligands from the solid supports in said library, and testing said ligands in an assay which identifies compounds having the desired characteristic.

12. A method of Claim 11 further comprising determining the structure of any ligand so identified.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/08882

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : G01N 33/53, 33/531; C07K 17/00; C07H 1/00

US CL : 436/518, 501; 435/6, 7.1; 530/333,334; 514/1; 536/18.5, 25.3

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 436/518, 501; 435/6, 7.1; 530/333,334; 514/1; 536/18.5, 25.3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, STN

search terms: combinatorial library, solid-phase synthesis

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y -- A	WO, A, 92/00091 (BIOLIGAND, INC.) 09 January 1992, see pages 6-7.	1, 2, 7-12 ----- 3, 5, 6 ----- 1-3, 5-12
Y -- A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCE, VOLUME 90, issued August 1993, DEWITT ET AL., "DIVERSOMERS": AN APPROACH TO NONPEPTIDE, NONOLIGOMERIC CHEMICAL DIVERSITY", pages 6909-6913, see entire document.	3, 4 ----- 3, 4



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be part of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Z" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

19 SEPTEMBER 1995

Date of mailing of the international search report

13 OCT 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

LORA M. GREEN

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/08882

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y -- A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCE, VOLUME 90, issued November 1993, NEEDELS ET AL., "GENERATION AND SCREENING OF AN OLIGONUCLEOTIDE-ENCODED SYNTHETIC PEPTIDE LIBRARY", pages 10700-10704, see Abstract.	5 -- 5
Y -- A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCE, VOLUME 90, issued December 1993, OHLMEYER ET AL., "COMPLEX SYNTHETIC CHEMICAL LIBRARIES INDEXED WITH MOLECULAR TAGS", pages 10922-10926, see Abstract.	6 -- 6

THIS PAGE BLANK (USPTO)